

γ -ray flux from Dark Matter Annihilation in Galactic Caustics

Lidia Pieri^{‡§} and Enzo Branchini[‡]

[‡] Department of Physics, Stockholm University, AlbaNova University Center
SE-10691 Stockholm, Sweden

[‡] Department of Physics, Università di Roma Tre
Via della Vasca Navale 84
I-00146 Rome, Italy

Abstract. In the frame of indirect dark matter searches we investigate the flux of high-energy γ -ray photons produced by annihilation of dark matter in caustics within our Galaxy under the hypothesis that the bulk of dark matter is composed of the lightest supersymmetric particles. Unfortunately, the detection of the caustics annihilation signal with currently available instruments is rather challenging. Indeed, with realistic assumptions concerning particle physics and cosmology, the γ -ray signal from caustics is below the detection threshold of both Čerenkov telescopes and satellite-borne experiments. Nevertheless, we find that this signal is more prominent than that expected if annihilation only occurs in the smoothed Galactic halo, with the possible exception of a $\sim 15^\circ$ circle around the Galactic center if the mass density profile of our Galaxy exhibits a sharp cusp there. We show that the angular distribution of this γ -ray flux changes significantly if DM annihilation preferentially occurs within virialized sub-halos populating our Galaxy rather than in caustics.

PACS numbers: 95.35.+d, 98.35.Gi, 98.35.Jk, 98.62.Gq, 11.30.Pb, 98.56.-p, 98.70.Rz

1. Introduction

Unveiling the nature and distribution of the dark matter (DM) is a fundamental goal of both theoretical and experimental physics today. The latest observations [1] allow $\sim 23\%$ of the matter in the universe to be of unknown, non-baryonic nature. One of the strongest candidates for the DM is a weakly interacting massive particle that was non-relativistic at the epoch of decoupling from the rest of the universe, as required by the currently popular cold dark matter (CDM) cosmological model. A good DM candidate is represented by the lightest supersymmetric particle (LSP) that in most particle

[§] e-mail: lidia@physto.se

physics scenarios is the neutralino χ . This particle is a spin- $\frac{1}{2}$ Majorana fermion constituted of a linear combination of the neutral gauge bosons and neutral Higgs doublet spartners, $\chi = a\tilde{B} + b\tilde{W}_3 + c\tilde{H}_1^0 + d\tilde{H}_2^0$. In the popular SUGRA or SUGRA-like models, where gaugino-universality is required, its mass is constrained by LEP limits to be greater than ~ 50 GeV [2] while theoretical considerations place an upper limit of a few hundred GeV - few TeV [3, 4] to the neutralino mass. If R -parity is conserved, neutralinos can change their cosmological abundance only through annihilation. Hence, high density regions constitute preferential locations for indirect DM searches.

Most of the recent works have assumed that the bulk of DM is constituted of neutralinos and looked for a possible detection of γ -ray annihilation flux with appropriate detectors. Phenomenological approaches to DM searches have focused mainly on the Galactic Center (GC) that has been observed by the EGRET satellite in the energy range 100 MeV - 10 GeV [5] and, more recently, by the ground-based Čerenkov telescopes WHIPPLE [6], CANGAROO-II [7] and HESS [8] with an energy threshold of hundreds of GeV. These data have shown an enhanced γ -ray flux along the the GC direction that has been variously interpreted and, although puzzling, does not yet constitute a firm evidence of indirect DM detection. New data from VERITAS [9] and MAGIC [10] Čerenkov telescopes, as well as from the soon-to-come satellite experiment GLAST [11], will certainly help clarifying this issue.

If dark relics are stable and their annihilation cross-section is non-negligible, then regions of enhanced density constitute preferential locations for the annihilation of DM particles. Our aim is to quantify the DM annihilation rate and to predict the associated γ -ray flux occurring within those singularities that arise in irrotational flows in a cold, collisionless medium during the process of structure formation, usually referred to as “caustics” because of their analogy with caustics in geometrical optics [12]. Although the presence of DM caustics is a general prediction of CDM cosmologies, model details depend on the characteristics of the DM candidate and on our ability in following the build up of caustics in the very nonlinear dynamical regime [13, 14, 15, 16]. In this work we will assume that the DM is composed of neutralinos and that the shape and position of Galactic caustics can be modeled as in [15].

The rationale behind this work is to predict the annihilation signal from DM caustics within our Galaxy, characterize its spatial signature and evaluate whether currently available or next-generation γ -ray detectors will be able to detect this signal and dis-

criminate it from the γ -ray Galactic background. Our results will be compared with those obtained from a previous similar analysis performed by [17, 18] that, however, considered the case of a single, nearby caustic of non-negligible transverse size.

The centers of virialized DM halos also constitute possible sites for annihilation of DM particles if a singularity in the mass density field called “cusp” develops there as a result of the gravitational instability process. Theoretical considerations and numerical experiments have neither been able to set strong constraints on the shape of the density profile near the halo center [19, 20, 21, 22, 23, 24, 25] nor to predict the abundance of a population of dark galactic sub-halos and their survival time within the Galactic halo [26, 27, 28, 29, 30]. On the other hand, current observations such as the rotation curves in the central galactic regions [31, 32, 33], Planetary Nebulae kinematics [34], microlensing events toward the Galactic bulge [35] and spectroscopic gravitational lensing [36] do not allow yet to discriminate among the various scenarios, although they seem to favor a very shallow, or cored density profile. Nevertheless, many authors have considered the possibility of detecting the products of DM annihilation occurring within Galactic, extragalactic and sub-galactic DM halos, taking into account both observational and theoretical constraints [37, 38, 25, 39, 40, 41, 42].

As neither theoretical arguments nor numerical experiments have been able yet to construct self consistent dynamical models that account for both DM caustics and virialized substructures populating a host DM matter halo it is difficult to quantitative compare the annihilation flux produced within either structures. Nevertheless, as we discuss in this work, qualitative differences between the angular distribution of the signals exist that may help in discriminating among these two scenarios.

The layout of the paper is as follows. In Section 2 we describe the expected photon flux from neutralino annihilation and introduce the detectability parameters for Čerenkov telescopes and satellite-borne experiments. In Section 3 we study the effect of a population of DM caustics on the expected γ -ray flux. Section 4 is devoted to investigating the possibility of detecting high energy photons produced in DM caustics. In Section 5 we summarize and discuss our main conclusions.

2. Detecting γ -rays from neutralino annihilation

Following the prescriptions given in [37], we factorize the diffuse photon flux from neutralino annihilation as

$$\frac{d\Phi_\gamma}{dE_\gamma}(E_\gamma, \psi, \Delta\Omega) = \frac{d\Phi^{\text{SUSY}}}{dE_\gamma}(E_\gamma) \times \Phi^{\text{cosmo}}(\psi, \Delta\Omega), \quad (1)$$

where ψ is the angle of view from the GC and $\Delta\Omega$ is the solid angle of observation for the detector. We will not consider here the γ -line emission. Its branching ratio is in fact as small as 10^{-3} and it produces a very small photon flux which we will neglect. In Eq. 1 the particle physics is embedded in the term:

$$\frac{d\Phi^{\text{SUSY}}}{dE_\gamma}(E_\gamma) = \frac{1}{4\pi} \frac{\langle\sigma_{\text{ann}}v\rangle}{2m_\chi^2} \cdot \sum_f \frac{dn_\gamma^f}{dE_\gamma} b_f \quad (2)$$

where $\langle\sigma_{\text{ann}}v\rangle$ is the thermally-averaged neutralino self-annihilation cross-section times the relative velocity of the two annihilating particles, dn_γ^f/dE_γ is the differential photon spectrum for a given annihilation final state f , whose analytical expression is given in [37], b_f is the branching ratio of annihilation into f and m_χ is the neutralino mass.

Astrophysics and cosmology affect the last factor of Eq. 1, that represents the line-of-sight integral:

$$\Phi^{\text{cosmo}}(\psi, \Delta\Omega) = \int_{\Delta\Omega} d\Omega' \int_{\text{l.o.s}} \rho_\chi^2(r(\lambda, \psi')) d\lambda(r, \psi'), \quad (3)$$

where $\rho_\chi(r)$ is the neutralino density profile, r is the distance from the GC and is equal to $r = \sqrt{\lambda^2 + R_\odot^2 - 2\lambda R_\odot \cos\psi}$, λ is the distance from the observer and $R_\odot$ is the distance of the Sun from the GC.

For distant, extragalactic objects, Eq. 3 reads

$$\Phi^{\text{cosmo}}(\psi, \Delta\Omega) = \frac{1}{d^2} \int_0^{\min[R_G, r_{\text{max}}(\Delta\Omega)]} 4\pi r^2 \rho_\chi^2(r) dr, \quad (4)$$

where d is the distance of the object from the observer, R_G is its virial radius and $r_{\text{max}}(\Delta\Omega)$ is the radius seen within the angular acceptance of the detector.

As mentioned before, the mass distribution near the center of the DM halo is poorly constrained. High resolution numerical experiments have shown that a cuspy density profile develops near the halo centers with an inner slope that is bracketed between the NFW profile (hereafter NFW97) [19]

$$\rho_\chi^{\text{NFW97}} = \frac{\rho_s^{\text{NFW97}}}{(r/r_s^{\text{NFW97}})(1 + r/r_s^{\text{NFW97}})^2} \quad (5)$$

and the steeper Moore et al. profile (M99) [20]:

$$\rho_\chi^{\text{M99}} = \frac{\rho_s^{\text{M99}}}{(r/r_s^{\text{M99}})^{1.5} [1 + (r/r_s^{\text{M99}})^{1.5}]}. \quad (6)$$

Scale quantities are fixed by the observations (virial mass or peak rotation velocity of the halo) and by the estimation of the concentration parameter $c = r_{\text{vir}}/r_s$, where we define the virial radius r_{vir} as the radius within which the halo average density is 200 times the critical density of the universe. The parameters c_{NFW97} and $c_{\text{M99}} = 0.64 c_{\text{NFW97}}$ have been computed according to [43] assuming a CDM universe.

As discussed in Section 1, the existence of a central cusp has not been confirmed by recent astronomical observations that, instead, seem to favor the case of a central core of constant density. Cored density profile would significantly depress the annihilation rate in the central region [44]. However, since using a cored rather than a cuspy density profile would not significantly affect our flux predictions for $\psi > 15^\circ$, where the caustics signal is expected to be more prominent, in this work we only consider the M99 and NFW97 model density profiles.

Fig.1 shows the factor $\Phi^{\text{cosmo}}(\psi)$ of Eq. 3 for our Galaxy and the three more prominent galaxies within a distance of 1.5 Mpc, computed for a solid angle of 10^{-5} sr (left panel) and 10^{-3} sr (right panel) assuming both a NFW (filled dots) and a Moore profile (filled triangles). We would like to remark here that the use of a cored profile would substantially reduce the expected fluxes.

Divergences in the l.o.s. integral are avoided by forcing a constant density core of radius 10^{-8} kpc corresponding to the distance at which the self-annihilation rate equals the dynamical time. Several astrophysical effects can increase the size of this core radius, leading to a significant decrease in the annihilation rate and expected flux [40, 37]. As already pointed out by [40], these external galaxies shine above the Galactic foreground as clearly seen in both panels of Fig.1.

The “supersymmetric factor” $\frac{d\Phi^{\text{SUSY}}}{dE_\gamma}$ in Eq. 2 has been investigated extensively by [37]. In our calculations we assume $\Phi^{\text{SUSY}} = 10^{-32} \text{ cm}^3 \text{ s}^{-1} \text{ GeV}^{-2} \text{ sr}^{-1}$ which represents a reasonable though somewhat optimistic choice among the values allowed by supersymmetric models.

2.1. Experimental detectability

One of the aims of this work is to study the experimental detectability of DM γ -ray sources with both Čerenkov telescopes and satellite-borne experiments. Following [37, 40], we explore the detectability of the diffuse photon flux from DM annihilation by comparing the number n_γ of expected γ events from neutralino annihilation with the




Figure 1. “Cosmological factor” $\Phi^{\text{cosmo}}(\psi)$ for the expected γ -ray flux from the most prominent external galaxies and for the Milky Way foreground within a solid angle $\Delta\Omega = 10^{-5}\text{sr}$ (left panel) or $\Delta\Omega = 10^{-3}\text{sr}$ (right panel), for a M99 (solid line and triangles) and a NFW97 (dotted line and circles) profile. A small central core region of radius 10^{-8}kpc has been assumed.

fluctuation of background events n_{bkg} :

$$\sigma \equiv \frac{n_{\gamma}}{\sqrt{n_{\text{bkg}}}} = \frac{\sqrt{T_{\delta}} \epsilon_{\Delta\Omega}}{\sqrt{\Delta\Omega}} \frac{\int A_{\gamma}^{\text{eff}} [d\phi_{\gamma}^{\text{DM}}/dEd\Omega] dEd\Omega}{\sqrt{\int \sum_{\text{bkg}} A_{\text{bkg}}^{\text{eff}} [d\phi_{\text{bkg}}/dEd\Omega] dEd\Omega}}. \quad (7)$$

To estimate the experimental sensitivity, σ , we choose the subsequent, rather typical set of parameters. For the effective observation time, T_{δ} , we consider a 20-day pointing for Čerenkov telescope and a 30-day observation for the satellite-borne experiment (for a Čerenkov telescope, only periods corresponding to a source zenith angle $\theta \leq 60^{\circ}$ are considered). The fraction of signal events $\epsilon_{\Delta\Omega}$ is set equal to 0.7 within the optimal solid angle $\Delta\Omega = 10^{-5}\text{sr}$, corresponding to an angular resolution of the instrument of 0.1° . The effective detection areas for electromagnetic and hadronic induced showers are considered to be independent from the value of the particle energy and incident angle and are set equal to $A_{\text{Čerenkov}}^{\text{eff}} = 4 \times 10^8 \text{cm}^2$ and $A_{\text{satellite}}^{\text{eff}} = 10^4 \text{cm}^2$, respectively. The identification efficiency ϵ is of crucial importance to reduce the physical background level. A Čerenkov telescope has a typical identification efficiency $\epsilon_{\text{e.m.}} \sim 99\%$ for electromagnetic induced (primary γ or electrons) showers and $\epsilon_{\text{had}} \sim 99\%$ for hadronic showers. A satellite experiment has an identification efficiency for charged particles of $\epsilon_{\text{charged}} \sim 99.997\%$ and of $\epsilon_{\text{neutral}} \sim 90\%$ for photons, due to the backlash of high energy photons [45]. Our expected signal is compared to the presence of proton [46], electron

[47] and photon [48, 49] backgrounds.

3. The caustics scenario

The caustics phenomenon has been extensively studied by [50]. Here we briefly outline the main characteristics of the DM caustics that are relevant for our work.

Collisionless dark matter particles move in the universe on a thin sheet in the velocity-position phase space, whose thickness is determined by their small primordial velocity dispersion. For a neutralino such a velocity has been calculated, after the equivalence time and before the onset of galaxy formation, to be as small as $\delta v_\chi(t) \sim 10^{-11}c (\text{GeV}/m_\chi)^{1/2} (t_0/t)^{2/3}$, where t_0 is the age of the universe. Such a sheet winds up where an overdensity grows non-linearly (i.e. when a galaxy forms) and the particles on the sheet fall onto the corresponding overdensity. The distance from the center of the overdensity at which the folding occurs has been estimated to be of the order of 1 Mpc for a galaxy like the Milky Way.

The following period of continuous cross-infalls lasts until inhomogeneities diffuse the process and prevent the formation of structures. In this period, a set of caustic surfaces which wrap around the galaxy is formed where the phase space folds. These surfaces define areas of enhanced density located at the boundary between regions characterized by different numbers of flows of particles. Namely, in the internal region of the caustic surface there is an odd number n of streams of particles compared to the $n-2$ flows in the external region. The non-zero velocity dispersion prevents the mass density to diverge at the caustics surface. More precisely, the density at the caustics is smeared off on distances $\delta x_\chi \sim 3 \cdot 10^{15} \text{ cm } (\text{GeV}/m_\chi)^{1/2}$, corresponding to about 10^{-4} pc for a TeV-mass neutralino. The final outcome of this process of caustics formation is the onset of a series of caustics with different characteristics. A number of *outer* caustics form, that appear as topological spheres surrounding the galaxy, as well as *inner* caustics, which look like tubes lying in the Galactic plane, composed of the particles which carry the most angular momentum when they are at their distance of closest approach to the Galactic Center. At the intersection between the Galactic plane and a perpendicular plane passing through the GC, these caustic tubes have a triangular shaped cross-section with three density cusps at the vertices of the triangle (the so-called tricusp), one on the Galactic plane and the other two located at the extremes of a segment perpendicular to it.

3.1. Density distribution near caustics

The particle density at the caustics can be calculated analytically [13]. In proximity of the cusps of the triangle the density profile d^{cau} scales like r^{-1} while near both the surface of the outer caustics and the side of the triangle it scales like $r^{-1/2}$, with r being the distance from the caustics. A set of simplifying assumptions is usually made to model the density distribution in these regions.

The claimed observational evidence of a Galactic tube characterized by a triangle with side of 100 pc [13] is at variance with the theoretical prediction of 1 kpc for the side of the triangle of the nearest inner caustics, suggesting that neither observations nor theory are currently able to set precise constraints on the size of caustic tubes. Therefore, we make the assumption that for all practical purposes the caustic triangles are of negligible sizes, i.e that the inner caustics reduce to caustic rings, or lines, on the Galactic plane. Such an approximation is valid [13] as long as the distance of the observer from the caustic ring is of the same order as the ring radius. This means that, assuming a galactocentric distance of 8.5 kpc, caustic rings with radii smaller than about 4-5 kpc can be considered as lines. Yet, the approximation still holds for caustic rings which are closer to the observer, provided that their transverse dimensions are smaller than the distances from the observer which, in this case, are smaller than the ring radius. In the model proposed by [15] the nearest caustics is characterized by a radius of 7.8 kpc and transverse dimensions of ~ 200 pc. When compared to the galactocentric distance, we have that $200 < 700 < 7.800$ pc, i.e. both the above conditions are satisfied and the approximation is valid. The previous relation is also valid for the remaining caustics that have similar transverse dimension but have larger radii. To compute the profile of the γ -ray emission from neutralino annihilation in the caustics we furthermore assume that the turnaround sphere is initially rigidly rotating and that the velocity of each particle in the flow is constant. Under these hypotheses we obtain a simple expression for the density distribution near the caustics.

For the the n -th outer caustics we have [14, 50]:

$$d_n(r) = 2 \frac{dM}{d\Omega dt} \bigg|_n \frac{1}{r^2 v_{\text{rot}} \sqrt{2 \ln \left(\frac{R_n}{r} \right)}}, \quad (8)$$

where r is the distance from the GC, v_{rot} is the rotational velocity of the galaxy, R_n is the radius of the outer caustics and $dM/d\Omega dt$ is the rate at which mass falls in per unit time and unit solid angle, that can be estimated using the self-similar infall model (see [14] for detailed references).

The density profile near the m -th caustic ring is given by [14, 50]:

$$d(a_m; \rho, z) \simeq \left. \frac{dM}{d\Omega dt} \right|_m \frac{2}{v} \frac{1}{\sqrt{(r^2 - a_m^2)^2 + 4a_m^2 z^2}} \quad (9)$$

where (ρ, z, θ) are cylindrical coordinates with respect to the GC, a_m is the caustics ring radius [15], $r = \sqrt{\rho^2 + z^2}$ and v is the velocity amplitude of the DM particles at the caustics position.

3.2. Expected γ -ray flux from caustics

Eqs. 8 and 9 can be used to compute the expected γ -ray flux from the caustics in our Galaxy. The resulting value of the “cosmological factor” $\Phi^{\text{cosmo}}(\psi)$ computed at the Galactic plane is shown in Fig. 2 for a solid angle of 10^{-5} sr (left panel) and 10^{-3} sr (right panel). The expected flux from caustics is obtained by integrating Eq. 3 over a region in which the mass density, given by either Eqs. 8 or 9, exceeds that of the underlying smooth Galactic halo. The mass within caustics, hence the amplitude of the annihilation signal, only depends on the virial mass of the host halo (or its rotational velocity). Indeed, in the self-similar infall model used to derive Eqs. 8 and 9, the value of v_{rot} determines both the rate of infall mass and caustics’ distances from the center of the halo.

The contribution from the outer caustics is negligible on the scale of the plots. A most remarkable spatial modulation signature is present which is due to the presence of the inner caustics rings ($n \geq 5$) with an associated signal well above the Galactic foreground level for both the M99 (straight line) and the NFW97 (dashed line) model density profiles.

The contribution of the caustics to the annihilation signal is negligible for extragalactic objects. In Fig. 2 we show the case of M31 (filled triangle and starred symbol), for which the effect of concentrating part of the halo mass in caustics actually decreases the flux. This derives from having placed a sizable fraction of the M31 mass in the outer caustics falling outside the viewing angle of the instrument.

Fig. 3 shows the ratio $(\phi^{MW} + \phi^{\text{cau}})/\phi^{MW}$, which defines the enhancement of the γ -ray signal due to the presence of the caustics rings, for a M99 (straight line) and a NFW97 (dotted line) profile. As expected, no enhancement is found in the direction of the GC, where the halo emissivity is very large, while DM annihilation in the nearest caustics ring results in an enhancement factor of about 600 along the direction $\psi = 60^\circ$. The spatial signature of the γ -ray flux expected in presence of Galactic caustics is indeed rather prominent.




Figure 2. “Cosmological factor” for the expected γ -ray emission from neutralino annihilation in the Galactic plane due to the presence of caustics rings for an angular resolution of 0.1 degrees corresponding to 10^{-5} sr (left panel) and 1 degree corresponding to 10^{-3} sr (right panel). The effect of the presence of caustics in the M31 galaxy is shown as well (symbols).




Figure 3. Enhancement of the γ -ray emission from neutralino annihilation in the Galactic plane due to the presence of caustics, for a solid angle of 10^{-5} sr.

3.3. Ring vs. Tricusp

As we have anticipated, neither theory nor observations are able to set strong constraints on the actual size of the caustics that ranges from ~ 100 pc to ~ 1.0 kpc. It is therefore worth investigating how our model predictions change when dropping our hypothesis of a ring caustics with negligible transverse size. This is precisely the case studied by [17] that considered the case of the nearest inner caustics (with $n=5$) located at 7.8 kpc from the Galactic Center with transverse dimensions of 1.2 kpc. With this particular geometry the Earth, placed at 8.5 kpc from the GC, is located well inside the tricusp.

The model [17] also assumes that $m_\chi = 100$ GeV instead of our fiducial value of 1 TeV, resulting in a larger smearing distance δx_χ and in a smaller cut-off on the caustic density profile of ~ 800 GeV cm $^{-3}$. Decreasing the density cutoff reduces the expected annihilation flux. The reduction factor weakly depends on the geometry of the problem. In the case of a single caustic ring with negligible size, the value of $\Phi_{\text{cosmo}}^{\text{MAX}}$ decreases by a factor of ~ 2 , where $\Phi_{\text{cosmo}}^{\text{MAX}}$ is the maximum of $\Phi_{\text{cosmo}}(\psi, \Delta\Omega = 10^{-5} \text{ sr}^{-1})$. In the case of a tricusp geometry considered by [17] the effect is very similar and $\Phi_{\text{cosmo}}^{\text{MAX}}$ decreases by a factor of ~ 3.3 .

Varying the geometry while keeping m_χ constant has a much more dramatic impact. We have found that with $m_\chi = 100$ GeV the value of $\Phi_{\text{cosmo}}^{\text{MAX}}$ for a tricusp is ~ 100 smaller than that expected from a caustic ring with negligible transverse size. This effect is hardly surprising as in the tricusp scenario, and unlike in the ring case, only a small portion of the caustics is seen within the angle of view $\Delta\Omega$.

Overall, the value of $\Phi_{\text{cosmo}}^{\text{MAX}}$ for the caustics $n = 5$ computed as in Section 3.2 is expected to be ~ 100 times larger than that of [17]. Note, however, that discrepancy in the flux predictions of the two models are less severe. This derives from the fact that our value of $\Phi^{\text{SUSY}} = 10^{-32} \text{ cm}^3 \text{ s}^{-1} \text{ GeV}^{-2} \text{ sr}^{-1}$ is ~ 1000 smaller than that assumed by [17] which, however, requires very optimistic hypotheses about the underlying particle physics.

4. Possibility of γ -ray flux detection

To investigate the possibility of detecting the flux of γ -ray photons produced by neutralinos annihilating in DM caustics, we compute the sensitivity of both a Čerenkov ground-based telescope and a satellite-borne experiment to the annihilation fluxes shown in Fig. 2. The parameters used to compute the experimental sensitivity (Eq. 7) are specified in Section 2.1 and represent realistic observations that could be carried out by

next-generation detectors.

Fig. 4 shows the experimental sensitivity, σ , a function of the angle from the GC in the case of a Čerenkov telescope at the VERITAS site (left panel) and a GLAST-like satellite (right panel). The choice of the Čerenkov telescope site determines the observable region of the sky and thus may constitute a source. For instance the GC region, which is maybe the brightest spot in the γ -ray sky, cannot be observed at the VERITAS site considered in this work. It is worth stressing that, because of its narrow field of view, each point in the left panels of Fig. 4 represents a single, 20-day long observation with a Čerenkov telescope. In both panels the significance of the detection is proportional to the γ -ray flux and thus exhibits the same dependence from ψ found in Fig. 2.

Clearly, the significance detection level is very low, even in correspondence of the peaks of the experimental sensitivity. This means that γ -ray photons produced by DM annihilation in the caustics cannot be detected by currently planned experiments unless new theoretical input from particle physics could boost up the supersymmetric factor by a factor of 30, which seems rather implausible.

These pessimistic conclusions are not alleviated when taking into account for the large theoretical uncertainties in the caustics model. Indeed, as we have shown in Section 3.3, increasing the transverse dimension of the nearest caustics while keeping constant its distance from the GC and using the same value of m_χ , has the effect of reducing the expected annihilation signal by a factor of ~ 100 making it more challenging its experimental detection. Moreover, a further decrease of the annihilation signal is to be expected if dynamical disturbances within the Galactic halo are large enough to significantly shorten their survival time ([27]).

5. Conclusions

In the popular CDM cosmological model, high density sub-structures are expected to form via gravitational instability from small perturbations in the primordial density field of cold, collisionless particles. In this work we have investigated the annihilation of DM particles within caustics, that represent regions of very high density, and have estimated the possibility of detecting the γ -ray photons produced by annihilation events within our Galaxy. Our main assumptions concern the geometry of the inner caustics, that we have considered small enough to be described as a series of line rings lying in the Galactic plane at various distances from the GC, as well as the underlying particle physics, namely the value of the neutralino mass $m_\chi = 1 \text{ TeV}$ and of $\Phi^{\text{SUSY}} = 10^{-32} \text{ cm}^3 \text{ s}^{-1} \text{ GeV}^{-2} \text{ sr}^{-1}$.




Figure 4. Left panel: expected sensitivity of a Čerenkov telescope to γ -ray emission from neutralino annihilation in the Galactic plane due to the presence of caustic rings, at an energy threshold of 100 GeV for a solid angle of 10^{-5} sr. Right panel: the same as in the left panel calculated for a satellite-borne experiment and an energy threshold of 10 GeV.

Our main conclusion is that the expected annihilation signal from inner Galaxy caustics cannot be revealed with presently available or currently planned Čerenkov telescopes and satellite-borne detectors. This result takes into account the present uncertainties in modeling the caustics geometry and their spatial distribution, both of which systematically contribute in suppressing the expected annihilation signal. As varying m_χ does not change significantly the value of Φ^{cosmo} , the only possibility of observing an annihilation signal from DM caustics would be that of assuming a much larger value for Φ^{SUSY} . For example, [17, 18] found that a significant detection is indeed possible in one year of observation with GLAST if a very optimistic value of $\Phi^{\text{SUSY}} = 10^{-29} \text{ cm}^3 \text{ s}^{-1} \text{ GeV}^{-2} \text{ sr}^{-1}$ is assumed that, however, we regard as rather implausible.

Although the possibility of its detection is small, some additional considerations about the annihilation signal are worth being made. First of all, the annihilation flux from Galaxy caustics dominates over the smooth Galactic background for $\psi > 15^\circ$ and possibly also within this region if a constant density core, rather than a sharp cusp, is present in the GC. Moreover, as we have checked, the annihilation signal from DM caustics does not significantly contribute to the total annihilation signal expected from

extragalactic sources such as M31.

Finally, it is interesting to compare the annihilation flux in the caustics scenario with that expected from a population of virialized sub-galactic halos that has already been studied by [26, 27, 28, 29, 30]. Indeed, the annihilation signal produced in Galaxy caustics is expected to have an angular distribution across the sky that is very different from that produced in the smooth Galactic halo (Fig. 2) and from that emitted by a population of virialized sub-galactic halos. In Fig. 5 we show the enhancement factor within a solid angle of 10^{-5} sr due to the presence of sub-Galactic halos of masses $\geq 10^6 M_\odot$ containing 20% of the original halo mass and having the same NFW97 density profile as the Galactic halo. The enhancement factor obviously depends on the modeling of the sub-Galactic halos density profile, mass function and spatial distribution (e.g. the three histograms of Fig. 5 show the effect of using the different models of sub-halos tidal disruptions proposed by [40]). Moreover, we would like to point out again that the use of a cored subhalos' density profile instead of the proposed NFW97 would further decrease the expected fluxes. Yet, as we have verified, accounting for model uncertainties does not significantly affect the angular distribution of the annihilation signal, in the sense that the enhancement profile has a peak that is broader than that predicted by the caustics scenario with a maximum at larger angles from the GC ($\psi \sim 120^\circ$ instead of $\sim 60^\circ$). The enhancement level remains low too, which means that γ -ray emission from sub-Galactic halos will hardly be detected by future gamma-ray detectors, especially because observations by satellite-borne experiments like GLAST will be hampered by the very long exposure time required by the serendipitous nature of these γ -ray sources.

Based on these considerations, the GC remains the best place to look at for detecting γ -ray signal, as it is shown in Fig. 4, provided that a sharp DM cusp is present there. However, ground-based observatories located in the northern hemisphere could only observe the GC at a large zenith angle, hence introducing large observational errors. For Čerenkov telescopes located in the southern hemisphere, such as HESS and CANGAROO, as well as for satellite-borne detectors, the GC represents by far the brightest source of the γ -ray sky and will therefore give the best chance to detect a possible signature of neutralino annihilation. Much more sensitive detectors will be needed to investigate the nature of the DM sources responsible for such signal.

Acknowledgments

We warmly thank L. Bergström, C. Gunnarsson, S. Matarrese and P. Sikivie for very useful discussions and suggestions. LP acknowledges a Research Grant funded jointly by the Italian Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR) and by




Figure 5. Enhancement of the γ -ray emission from neutralino annihilation in the Galactic plane due to the presence of a population of sub-halos for a solid angle of 10^{-5} sr and a NFW97 profile. The corresponding average values of the enhancement factor are indicated by the three straight lines.

the University of Torino, and thanks the Department of Theoretical Physics of the University of Torino where this work has been partially carried on.

- [1] D. N. Spergel et al., *Astrophys. J. Suppl.* **148**, 175 (2003).
- [2] LEPSUSY Working Group, ALEPH, DELPHI, L3 and OPAL experiments, <http://lepsusy.web.cern.ch/lepsusy/Welcome.html>.
- [3] J. Ellis, et al., *Phys. Rev. D.* **62**, 075010 (2000).
- [4] G. Jungman, et al., *Phys. Rep.* **267**, 195 (1996).
- [5] S.D. Hunter *et al.*, *Astrophys. J.* **481**, 205 (1997).
- [6] K. Kosack *et al.*, *Astrophys. J.* **608**, L7 (2004).
- [7] K. Tsuchiya *et al.*, *Astrophys. J.* **606**, L115 (2004).
- [8] F. Aharonian *et al.*, astro-ph/0408145.
- [9] T. C. Weekes et al., in *Proc. of the 25th ICRC* **5**, 173 (1997).
- [10] C. Baixeras, et al., *Nucl.Phys.Proc.Suppl.*, **114**, 247, (2003).
- [11] A. Morselli, et al., in *Proc. of the 32nd Rencontres de Moriond*, (1997).
- [12] I. B. Zeldovich, A.V. Mamaev and S. F. Shandarin, *Soviet Physics - Uspekhi* **26**, 77 (1983).
- [13] P. Sikivie, *Phys. Rev. D* **60**, 063501 (1999).
- [14] V.K. Onemli, astro-ph/0401162.
- [15] P. Sikivie, in *Proceedings of the 2nd International Workshop on the Identification of Dark Matter*, Buxton, England, Sept. 7-11, 1998
- [16] A. Helmi, S.D.M. White and V. Springel, *Phys. Rev. D* **66**, 063502 (2002).
- [17] L. Bergström, J. Edsjö and C. Gunnarsson, *Phys. Rev. D* **63**, 083515 (2001).

- [18] C. Gunnarsson, *Neutralino Dark Matter and Caustic Ring Signals*, Diploma Thesis.
- [19] J. F. Navarro, et al., *Astrophys. J.*, **490**, 493 (1997).
- [20] B. Moore, et al., *Astrophys. J. Lett.* **524**, L19 (1999).
- [21] P. Ullio, H. Zhao and M. Kamionkowski, *Phys. Rev. D* **64**, 043504 (2001).
- [22] J. F. Navarro *et al.*, *MNRAS* **349**, 1039 (2004)
- [23] J. Diemand, B. Moore and J. Stadel, *MNRAS*, **352**, 535 (2004)
- [24] C. Power et al., *MNRAS.*, **338**, 14 (2003).
- [25] F. Stoehr, S.D.M. White, V. Springel, G. Tormen, N. Yoshida, *MNRAS* **345**, 1313 (2003).
- [26] E. Hayashi, et al., *Astrophys.J.* **584**, 541 (2003)
- [27] A. Helmi, S. White, V. Springel. *MNRAS.*, **339**, 834 (2003)
- [28] L. Gao, G. De Lucia, S. D. M. White and A. Jenkins, *MNRAS.*, **352**, L1 (2004)
- [29] J. Diemand, B. Moore and J. Stadel, *Nature* **433**, 389 (2005)
- [30] H. Zhao, J. Taylor J. Silk, D. Hooper, astro-ph/0502049
- [31] W. J. G. de Blok, A. Bosma and S. McGaugh, *MNRAS* **340**, 657 (2003).
- [32] R. A. Swaters, et al., *Astrophys.J.* **583**, 732 (2003).
- [33] F. Donato, G. Gentile, P. Salucci, *MNRAS*, **353**, L17 (2004);
A. Borriello and P. Salucci, *MNRAS* **323**, 295 (2001).
- [34] M.R. Merrifield, astro-ph/0412059
- [35] C. Afonso, et al., *A&A* **404**, 145, (2003).
- [36] R. B. Metcalf, L. A. Moustakas, A. J. Bunker and I. R. Parry *Astrophys. J.*, **607**, 43 (2004).
- [37] N. Fornengo, L. Pieri and S. Scopel, hep-ph/0407342.
- [38] V. Berezhinsky, A. Bottino, G. Mignola, *Phys. Lett.* **B325**, 136 (1994);
G. Jungman, M. Kamionkowski, *Phys. Rev. D* **51**, 3121 (1995);
L. Bergström, J. Edsjö, and P. Ullio, *Phys. Rev. D* **58**, 083507 (1998);
L. Bergström, J. Edsjö, P. Gondolo, and P. Ullio, *Phys. Rev. D* **59**, 043506 (1998);
L. Bergström, P. Ullio and J. Buckley, *Astropart. Phys.* **9**, 137 (1998);
P. Ullio and L. Bergström, *Nucl. Phys.* **B504**, 27 (1997); *Phys. Rev. D* **57**, 1962(1998);
C. Calcano-Roldan, B. Moore, *Phys. Rev. D* **62**, 123005 (2000);
G. Bertone, G. Sigl, J. Silk, *Mon. Not. Roy. Astron. Soc.* **326**, 799 (2001);
J. L. Feng, K. T. Matchev, F. Wilczek, *Phys. Rev. D* **63**, 045024 (2001);
A. Cesarini, F. Fucito, A. Lionetto, A. Morselli, P. Ullio, *Astropart. Phys.* **21**, 267 (2004);
R. Aloisio, P. Blasi, A. V. Olinto, *JCAP* **0405**, 007 (2004);
R. Aloisio, P. Blasi, A. V. Olinto, *Astrophys. J.* **601**, 47 (2004);
S. Peirani, R. Mohayaee, J. A. de Freitas Pacheco, *Phys. Rev. D* **70**, 043503 (2004);
Y. Mambrini and C. Munoz, hep-ph/0407158.
- [39] A. Bottino, F. Donato, N. Fornengo, S. Scopel, *Phys. Rev. D* **70**, 015005 (2004).
- [40] L. Pieri and E. Branchini, *Phys. Rev. D* **69**, 043512 (2004).
- [41] C. Calcano-Roldan and B. Moore, *Phys. Rev. D*, **62**, 1230 (2000).
- [42] P. Ullio, et al., *Phys. Rev. D* **66**, 123502 (2002);
J. Taylor and J. Silk, *MNRAS.*, **339**, 505 (2003);
L. Bergström, J. Edsjö, P. Ullio, *Phys. Rev. Lett.* **87**, 1301 (2001);
C. Tyler, *Phys. Rev. D* **66**, 023509 (2002);
E. A. Baltz, et al., *Phys. Rev. D* **61**, 023514 (2000);
A. Tasitsiomi, J. Gaskins, A. Olinto, astro-ph/0307375, to appear in *Astroparticle Physics*.
- [43] V. R. Eke, et al., *Astrophys.J.* **554**, 114 (2001).

- [44] N.W. Evans, F. Ferrer, S. Sarkar, Phys. Rev. D **69**, 123501 (2004);
- [45] A. Moiseev et al., Proc. of the 28th ICRC (2003).
- [46] T. Gaisser, et al., in Proc. of the 27th ICRC, (2001).
- [47] M. S. Longair, High Energy Astrophysics, Cambridge University Press, (1992).
- [48] L. Bergström, et al., Astroparticle Phys. **9**, 137 (1998).
- [49] P. Sreekumar, et al., ApJ **494**, 523, (1998).
- [50] E. Bertschinger, Ap. J. Suppl., **58**, 39 (1985);
J. R. Ipser and P. Sikivie, Phys. Lett. B, **291**, 288 (1992);
P. Sikivie, I. I. Tkachev and Y. Wang, Phys. Rev. Lett., **75**, 2911 (1995);
P. Sikivie, Phys. Lett. B, **432**, 139 (1998);
P. Sikivie, Phys. Rev. D, **60**, 063501 (1999).